




ELSEVIER

Contents lists available at ScienceDirect

Linear Algebra and its Applications

www.elsevier.com/locate/laa



On a conjecture of Ilmonen, Haukkanen and Merikoski concerning the smallest eigenvalues of certain GCD related matrices



Ercan Altınışık^{a,*}, Ali Keskin^a, Mehmet Yıldız^a,
Murat Demirbükten^b

^a Department of Mathematics, Faculty of Sciences, Gazi University, 06500
Teknikokullar – Ankara, Turkey

^b Department of Computer Science, Faculty of Engineering, İ. Doğramacı Bilkent
University, 06800 Bilkent – Ankara, Turkey

ARTICLE INFO

Article history:

Received 14 July 2015

Accepted 23 November 2015

Available online 15 December 2015

Submitted by R. Brualdi

MSC:

15A18

15A23

15B36

15B48

11B39

11C20

Keywords:

GCD matrix

(0, 1)-matrix

Positive matrix

Eigenvalue

Spectral radius

Fibonacci number

ABSTRACT

Let K_n be the set of all $n \times n$ lower triangular $(0, 1)$ -matrices with each diagonal element equal to 1, $L_n = \{YY^T : Y \in K_n\}$ and let c_n be the minimum of the smallest eigenvalue of YY^T as Y goes through K_n . The Ilmonen–Haukkanen–Merikoski conjecture (the IHM conjecture) states that c_n is equal to the smallest eigenvalue of $Y_0Y_0^T$, where $Y_0 \in K_n$ with $(Y_0)_{ij} = \frac{1-(-1)^{i+j}}{2}$ for $i > j$. In this paper, we present a proof of this conjecture. In our proof we use an inequality for spectral radii of nonnegative matrices.

© 2015 Elsevier Inc. All rights reserved.

* Corresponding author. Tel.: +90 312 202 1070.

E-mail addresses: ealtinisik@gazi.edu.tr (E. Altınışık), akeskin1729@gmail.com (A. Keskin), yildizm78@my.net (M. Yıldız), murdem91@gmail.com (M. Demirbükten).

1. Introduction

Let $S = \{x_1, x_2, \dots, x_n\}$ be a set of distinct positive integers, (x_i, x_j) denote the greatest common divisor of x_i and x_j and let ε be a positive real number. The $n \times n$ matrices $(S) = ((x_i, x_j))$ and $(S^\varepsilon) = ((x_i, x_j)^\varepsilon)$ are called the GCD matrix and the power GCD matrix on S , respectively. The LCM matrix and the power LCM matrix are similarly defined. In 1876, Smith [30] proved that if S is factor closed, then $\det(S) = \prod_{k=1}^n \varphi(x_k)$, where φ is Euler's totient. Since then many results on these matrices have been published in the literature, see e.g. [5,6,10,11,13,21].

An interesting and active area in the study of GCD type matrices is their eigenstructure. The first results on this subject were published in the papers [32,7,23] but the paper of Hong and Loewy [16] can be considered as the first paper on the study of the eigenvalues of GCD and related matrices due to the number theoretical aspect of the subject. Since their pioneering paper many results on the subject have been published in the literature, see e.g. [1,3,4,14,15,17,19,24,26–28]. In that paper, Hong and Loewy investigated the asymptotic behavior of the eigenvalues of power GCD matrices by using some tools of number theory. Beside their results on asymptotic behavior of these matrices, in the same paper Hong and Loewy introduced a constant c_n and used it to present a lower bound for the smallest eigenvalues of power GCD matrices. Let K_n be the set of all $n \times n$ lower triangular $(0, 1)$ -matrices with each diagonal element equal to 1 and let $L_n = \{YY^T : Y \in K_n\}$. They defined the numbers c_n depending only on n as follows:

$$c_n = \min_{Z \in L_n} \left\{ \mu_n^{(1)}(Z) : \mu_n^{(1)}(Z) \text{ is the smallest eigenvalue of } Z \right\}. \quad (1.1)$$

Then they proved that

$$\lambda_n^{(1)}((S^\varepsilon)) \geq c_n \cdot \min_{1 \leq i \leq n} \{J_\varepsilon(x_i)\},$$

where J_ε is Jordan's generalization of Euler's totient and $\lambda_n^{(1)}((S^\varepsilon))$ is the smallest eigenvalue of the power GCD matrix (S^ε) , see [16, Theorem 4.2].

In 2008, Ilmonen, Haukkanen and Merikoski [19] studied eigenvalues of meet and join matrices which are abstract generalizations of GCD and LCM matrices, respectively, and they generalized the above result concerning the numbers c_n for positive definite meet matrices defined on locally finite meet semilattices. They also obtained a similar result for join matrices and hence, in particular, for LCM matrices. In the same paper, in the light of their MATLAB calculations for $n = 2, 3, \dots, 7$, they presented an interesting conjecture about the constants c_n .

Conjecture 1.1 (The IHM conjecture). (See [19, Conjecture 7.1].) Let $Y_0 = (y_{ij}^0) \in K_n$ be defined by

$$y_{ij}^0 = \frac{1 - (-1)^{i+j}}{2} \quad (1.2)$$

if $i > j$. Then c_n is equal to the smallest eigenvalue of $Y_0 Y_0^T$.

We have further numerical evidence of the IHM conjecture as follows. Recently, the first, the second and the fourth author of the paper have investigated that the IHM conjecture holds for $n = 7$ and $n = 8$ with the help of a MATLAB code. Our MATLAB code running on a computer¹ has verified the truth of the IHM conjecture for $n = 7$ in 23 minutes and for $n = 8$ in 3.5 days. Since $|L_8| = 2^{28} = 268,435,456$ and $|L_9| = 2^{36} = 68,719,476,736$, it would take about 3 years to verify the IHM conjecture for $n = 9$ with the help of our MATLAB code. To overcome this difficulty about time, we write a different code in C programming language. We use Newton's identities (see [20]) to obtain the characteristic polynomial of a matrix Z in L_n and we calculate the smallest eigenvalue of Z by using Newton's method (see [31]) to shorten the running time. Indeed, our C code running on the same computer has verified the truth of the IHM conjecture in 30 minutes for $n = 8$ and in 7 days for $n = 9$. Thus, we have concluded that [Conjecture 1.1](#) holds for $n = 8$ and $n = 9$. This investigation has been presented by the first author of this paper in [2].

After obtaining enough numerical evidence that the IHM conjecture can be true, we get the motivation to find out a proof of it. The strategy of the proof is as follows. We prove that for any matrix Y in K_n , $|Y^{-1}| \leq |Y_0^{-1}|$, where the matrix Y_0 given by (1.2) and $|Y^{-1}|$ is the element-wise absolute value of the matrix Y^{-1} . Secondly, we show that $|Z^{-1}| \leq |(Y_0 Y_0^T)^{-1}|$ for all $Z \in L_n$. Then, by using an inequality for spectral radii of nonnegative matrices, we obtain a proof of the IHM conjecture. We conclude the paper with a conjecture on the uniqueness of the matrix Y_0 and a discussion about further studies on the constant c_n .

2. Properties of matrices in K_n

First we present a simple fact about a particular nilpotent $(0, 1)$ -matrix which we use in the course of our proofs. Here we give the proof of this fact though one can find in the literature.

Lemma 2.1. *Let $Y \in K_n$ and $N := Y - I$, where I is the $n \times n$ identity matrix. We denote by $(N^k)_{ij}$ the ij -entry of the positive integer k -th power of N . Then we have the following properties.*

- i) $(N^k)_{ij} = 0$ whenever $i - j < k$,
- ii) $Y^{-1} = I - N + N^2 - \cdots + (-1)^{n-1} N^{n-1}$.

¹ Intel Core i7-920 Quad Core 2.66 GHz 8 MB L Cache 24 GB DDR3 RAM.

Proof. Since N is a strictly lower triangular $(0, 1)$ -matrix, the proof of the first claim follows from the matrix multiplication. For the proof of the second claim, consider

$$I^n - (-N)^n = (I + N)(I - N + N^2 - \cdots + (-1)^{n-1}N^{n-1}).$$

Since $N^n = 0$ and $Y = I + N$, we have

$$Y^{-1} = I - N + N^2 - \cdots + (-1)^{n-1}N^{n-1}. \quad \square$$

Now we investigate the inverse of any matrix Y in K_n . In the following lemma we obtain a recurrence relation for the entries of the inverse of Y .

Lemma 2.2. *Let $Y \in K_n$ and $N := Y - I$. Denote $N = (n_{ij})$ and $Y^{-1} = (a_{ij})$. Then we have the following recurrence relation for a_{kl} :*

$$a_{kl} = \begin{cases} 0 & \text{if } k < l, \\ 1 & \text{if } k = l, \\ -\sum_{i=l}^{k-1} n_{ki}a_{il} & \text{if } k > l. \end{cases}$$

Proof. When we multiply both sides of the equality in Lemma 2.1(ii) from the left by $-N$, we have

$$-NY^{-1} = -N + N^2 - \cdots + (-1)^{n-1}N^{n-1} + (-1)^nN^n.$$

Since $N^n = 0$, one can easily obtain

$$I - NY^{-1} = Y^{-1}. \quad (2.1)$$

By Lemma 2.1, it is clear that $a_{kl} = 0$ if $k < l$ and $a_{kl} = 1$ if $k = l$. Now, from (2.1), we have

$$a_{kl} = -\sum_{i=1}^n n_{ki}a_{il}$$

for all $k > l$ and hence, by Lemma 2.1, we obtain

$$a_{kl} = -\sum_{i=l}^{k-1} n_{ki}a_{il}$$

for all $k > l$. \square

In the following theorem, we find an upper bound for each a_{ij} in terms of Fibonacci numbers which is a surprising result.

Theorem 2.1. Let $Y \in K_n$ and denote $Y^{-1} = (a_{ij})$. Then, for $1 \leq j < i \leq n$, we have $|a_{ij}| \leq f_{i-j}$, where f_{i-j} is the $(i-j)$ -th Fibonacci number.

Proof. Let $N = (n_{ij})$ be as in Lemma 2.2. Let $j = 1, 2, \dots, n-1$. We prove by induction on $t = 1, 2, \dots, n-j$ that $|a_{j+t,j}| \leq f_t$ for all $Y \in K_n$. By Lemma 2.2, we have $|a_{j+1,j}| = n_{j+1,j}$, where $n_{j+1,j}$ can be 0 or 1. Thus, $|a_{j+1,j}| \leq 1 = f_1$ for all $Y \in K_n$. Now, assume that for each $t = 1, 2, \dots, k-1$ we have $|a_{j+t,j}| \leq f_t$ for all $Y \in K_n$. We prove that $|a_{j+k,j}| \leq f_k$ for all $Y \in K_n$. By Lemma 2.2, we have $a_{j+k,j} = -\sum_{i=j}^{j+k-1} n_{j+k,i} a_{ij}$ for all $Y \in K_n$.

Case 1. Assume $n_{j+k,j+k-1} = 0$. Then $|a_{j+k,j}| = \left| \sum_{i=j}^{j+k-2} n_{j+k,i} a_{ij} \right|$. Also, by Lemma 2.2, $a_{j+k-1,j} = -\sum_{i=j}^{j+k-2} n_{j+k-1,i} a_{ij}$. Since both of $n_{j+k,i}$ and $n_{j+k-1,i}$ for each $i = j, j+1, \dots, j+k-2$ can arbitrarily be 0 or 1, it is clear that $a_{j+k,j}$ and $a_{j+k-1,j}$ go through the same values, as Y goes through the set K_n . Therefore, by the induction hypothesis, we obtain $|a_{j+k,j}| \leq f_{k-1} \leq f_k$ for all $Y \in K_n$.

Case 2. Assume $n_{j+k,j+k-1} = 1$.

Subcase i. Assume $n_{j+k-1,j+k-2} = 0$. By Lemma 2.2, we have

$$|a_{j+k,j}| \leq \left| \sum_{i=j}^{j+k-2} n_{j+k,i} a_{ij} \right| + |a_{j+k-1,j}|.$$

Also, by Lemma 2.2, it is clear that $a_{j+k-1,j} = -\sum_{i=j}^{j+k-2} n_{j+k-1,i} a_{ij}$. Since both of $n_{j+k,i}$ and $n_{j+k-1,i}$ for each $i = j, j+1, \dots, j+k-2$ can arbitrarily be 0 or 1, it is clear that $-\sum_{i=j}^{j+k-2} n_{j+k,i} a_{ij}$ and $a_{j+k-1,j}$ go through the same values, as Y goes through the set K_n . Thus, by the induction hypothesis, we obtain $\left| \sum_{i=j}^{j+k-2} n_{j+k,i} a_{ij} \right| \leq f_{k-1}$.

Beside this, by our assumption in Subcase i, $|a_{j+k-1,j}| = \left| \sum_{i=j}^{j+k-3} n_{j+k-1,i} a_{ij} \right|$. Since both of $n_{j+k-1,i}$ and $n_{j+k-2,i}$ for each $i = j, j+1, \dots, j+k-3$ can arbitrarily be 0 or 1, it is obvious that $\sum_{i=j}^{j+k-3} n_{j+k-1,i} a_{ij}$ and $a_{j+k-2,j}$ go through the same values, as Y goes through the set K_n . By the induction hypothesis, we obtain $|a_{j+k-1,j}| \leq f_{k-2}$ for all $Y \in K_n$. Thus, $|a_{j+k,j}| \leq f_{k-1} + f_{k-2} = f_k$ for all $Y \in K_n$.

Subcase ii. Assume $n_{j+k-1,j+k-2} = 1$. By Lemma 2.2, we have

$$|a_{j+k,j}| \leq \left| \sum_{i=j}^{j+k-3} n_{j+k,i} a_{ij} \right| + \left| \sum_{i=j+k-2}^{j+k-1} n_{j+k,i} a_{ij} \right|.$$

Since $\sum_{i=j}^{j+k-3} n_{j+k,i} a_{ij}$ and $a_{j+k-2,j}$ go through the same values, as Y goes through the set K_n , by the induction hypothesis, we have $\left| \sum_{i=j}^{j+k-3} n_{j+k,i} a_{ij} \right| \leq f_{k-2}$ for all $Y \in K_n$. In addition to this, since $n_{j+k-1,j+k-2} = 1$ we obtain

$$\sum_{i=j+k-2}^{j+k-1} n_{j+k,i} a_{ij} = (n_{j+k,j+k-2} - 1) a_{j+k-2,j} - \sum_{i=j}^{j+k-3} n_{j+k-1,i} a_{ij}.$$

Here each $(1 - n_{j+k,j+k-2}), n_{j+k-1,j}, \dots, n_{j+k-1,j+k-3}$ can arbitrarily be 0 or 1. Thus, $\sum_{i=j+k-2}^{j+k-1} n_{j+k,i} a_{ij}$ and $a_{j+k-1,j}$ go through the same values, as Y goes through the set K_n and hence, by the induction hypothesis, $\left| \sum_{i=j+k-2}^{j+k-1} n_{j+k,i} a_{ij} \right| \leq f_{k-1}$ for all $Y \in K_n$. Therefore, we obtain $|a_{j+k,j}| \leq f_{k-2} + f_{k-1} = f_k$ for all $Y \in K_n$.

The principle of induction completes the proof. \square

Let $A = (a_{ij}), B = (b_{ij}) \in M_n(\mathbb{R})$, that is, the set of all $n \times n$ real matrices. We write $A \geq 0$ if all $a_{ij} \geq 0$. Also, we write $A \geq B$ if $A - B \geq 0$. In addition to this, we define $|A| = (|a_{ij}|)$, that is, $|A|$ is the element-wise absolute value of A . The largest eigenvalue of A in modulus is denoted by $\rho(A)$ and called the spectral radius of A . Now we fix the notation for the rest of the paper.

Theorem 2.2. Let $Y_0 = (y_{ij}^0) \in K_n$ be as in (1.2) and let $Z_0 := Y_0 Y_0^T$. For all $Z \in L_n$, we have $|Z^{-1}| \leq |Z_0^{-1}|$.

Proof. Firstly we obtain the inverse of Y_0 . Let $N_0 = Y_0 - I$ and $N_0 = (m_{ij})$. Then

$$m_{ij} = \begin{cases} 0 & \text{if } i \leq j, \\ \frac{1 - (-1)^{i+j}}{2} & \text{otherwise.} \end{cases}$$

We claim that the inverse of Y_0 is the $n \times n$ matrix (c_{ij}) , where

$$c_{ij} = \begin{cases} 0 & \text{if } i < j, \\ 1 & \text{if } i = j, \\ (-1)^{i-j} f_{i-j} & \text{if } i > j. \end{cases}$$

Since $Y_0 \in K_n$ by Lemma 2.2, it is clear that $c_{ij} = 0$ if $i < j$ and $c_{ij} = 1$ if $i = j$. Also, by Lemma 2.2, we have

$$c_{ij} = - \sum_{k=j}^{i-1} m_{ik} c_{kj}$$

for $i > j$. Now we prove that $c_{ij} = (-1)^{i-j} f_{i-j}$ whenever $i > j$ by induction on $t = i - j$. For $t = 1$,

$$c_{j+1,j} = -m_{j+1,j} = -1 = -f_1.$$

Assume that $c_{j+t,j} = (-1)^t f_t$ for all $t = 1, 2, \dots, k-1$. Recall that

$$c_{j+k,j} = - \sum_{s=j}^{j+k-1} m_{j+k,s} c_{sj}.$$

By the induction hypothesis, if k is even then we have

$$c_{j+k,j} = \sum_{s=1}^{k/2} f_{2s-1} = f_k$$

and if k is odd then

$$c_{j+k,j} = -1 - \sum_{s=1}^{(k-1)/2} f_{2s} = -f_k.$$

Here the last equalities follow from the well-known Fibonacci identities, see [22, Theorem 5.2 and Corollary 5.1]. Thus, $c_{j+k,j} = (-1)^k f_k$.

Secondly, we calculate the inverse of Z_0 . Since $Z_0 = Y_0 Y_0^T$ and $Y_0^{-1} = (c_{ij})$, we have

$$(Z_0^{-1})_{ii} = \sum_{k=1}^n c_{ki}^2 = 1 + \sum_{k=i+1}^n f_{k-i}^2 \quad (2.2)$$

for all $i = 1, 2, \dots, n$. Now let $1 \leq i < j \leq n$. Then

$$\begin{aligned} (Z_0^{-1})_{ij} &= \sum_{t=1}^n c_{ti} c_{tj} \\ &= c_{ji} + \sum_{t=j+1}^n c_{ti} c_{tj} \\ &= (-1)^{j-i} f_{j-i} + \sum_{t=j+1}^n (-1)^{-i-j} f_{t-i} f_{t-j} \\ &= (-1)^{j-i} (f_{j-i} + \sum_{t=j+1}^n f_{t-i} f_{t-j}). \end{aligned} \quad (2.3)$$

Since Z_0^{-1} is symmetric, for all $1 \leq j < i \leq n$,

$$(Z_0^{-1})_{ij} = (-1)^{i-j} (f_{i-j} + \sum_{t=i+1}^n f_{t-i} f_{t-j}). \quad (2.4)$$

Now we prove the claim of the theorem. For each $Z \in L_n$, there exists a matrix Y in K_n such that $Z = Y Y^T$. Let $Y^{-1} = (a_{ij})$. Then, by Lemma 2.2 and Theorem 2.1, we have

$$\begin{aligned}
|(Z^{-1})_{ii}| &= \left| \sum_{k=1}^n a_{ki}^2 \right| \\
&= \sum_{k=1}^n |a_{ki}|^2 \\
&= \sum_{k=i}^n |a_{ki}|^2 \\
&\leq 1 + \sum_{k=i+1}^n f_{k-i}^2 \\
&= |(Z_0^{-1})_{ii}|
\end{aligned}$$

for all $i = 1, 2, \dots, n$. Let $j > i$. By [Lemma 2.2](#) and [Theorem 2.1](#), we have

$$\begin{aligned}
|(Z^{-1})_{ij}| &= \left| \sum_{t=1}^n a_{ti} a_{tj} \right| \\
&\leq \sum_{t=1}^n |a_{ti}| |a_{tj}| \\
&= |a_{ji}| + \sum_{t=j+1}^n |a_{ti}| |a_{tj}| \\
&\leq f_{j-i} + \sum_{t=j+1}^n f_{t-i} f_{t-j} \\
&= |(Z_0^{-1})_{ij}|.
\end{aligned}$$

Finally, since Z^{-1} and Z_0^{-1} are symmetric, $|Z^{-1}| \leq |Z_0^{-1}|$ for all $Z \in L_n$. \square

3. Proof of [Conjecture 1.1](#)

The following lemma is crucial in the proof of [Conjecture 1.1](#).

Lemma 3.1. (See [\[18, Theorem 8.1.18\]](#).) Let $A, B \in M_n(\mathbb{R})$. If $|A| \leq B$, then $\rho(A) \leq \rho(|A|) \leq \rho(B)$.

Proof of the conjecture. Let Z_0 be as in [Theorem 2.2](#). First we prove that the matrices Z_0^{-1} and $|Z_0^{-1}|$ have the same characteristic polynomial. By the definition of the trace of a square matrix, it is clear that

$$\text{tr}((Z_0^{-1})^k) = \sum_{i_1, \dots, i_k=1}^n (Z_0^{-1})_{i_1 i_2} \cdots (Z_0^{-1})_{i_{k-1} i_k} (Z_0^{-1})_{i_k i_1}$$

for each $k = 1, 2, \dots, n$. Also, from formulae (2.2)–(2.4) in the proof of Theorem 2.2, one can easily show that $\text{sgn}(Z_0^{-1})_{ij} = (-1)^{i-j}$ for all $1 \leq i, j \leq n$. Thus, we have

$$\text{sgn}((Z_0^{-1})_{i_1 i_2} \dots (Z_0^{-1})_{i_{k-1} i_k} (Z_0^{-1})_{i_k i_1}) = 1$$

and hence $\text{tr}(|Z_0^{-1}|^k) = \text{tr}((Z_0^{-1})^k)$. By Newton's identities [20], we obtain that Z_0^{-1} and $|Z_0^{-1}|$ have the same characteristic polynomial. Thus, $\rho(|Z_0^{-1}|) = \rho(Z_0^{-1})$. From Theorem 2.2 and Lemma 3.1, now we obtain

$$\rho(Z^{-1}) \leq \rho(|Z^{-1}|) \leq \rho(|Z_0^{-1}|) = \rho(Z_0^{-1}),$$

for all Z in L_n . Since all Z in L_n are positive definite, the smallest eigenvalue of Z_0 is less than or equal to the smallest eigenvalue of Z for all Z in L_n . \square

Finally, in our investigation [2], we cannot find any matrix Y other than Y_0 in K_n such that c_n is equal to the smallest eigenvalue of YY^T for each $n = 2, 3, \dots, 9$. After this observation we can present the following conjecture.

Conjecture 3.1. *Let $n \geq 2$. There is a unique matrix Y in K_n such that c_n is equal to the smallest eigenvalue of YY^T . In other words, if the smallest eigenvalue of YY^T is equal to c_n then $Y = Y_0$, where Y_0 is defined by (1.2).*

4. Lower bounds for c_n

In the literature there are not so many results on estimating the value of c_n . Recently, Mattila [24] has presented a lower bound for c_n . Indeed, he proved that c_n is bounded below by $(\frac{6}{n^4+2n^3+2n^2+n})^{\frac{n-1}{2}}$. Then he showed that this lower bound can be replaced with $(\frac{48}{n^4+56n^2+48n})^{\frac{n-1}{2}}$ when n is even, and $(\frac{48}{n^4+50n^2+48n-51})^{\frac{n-1}{2}}$ when n is odd. Recently, beside Mattila's results, Altınışık and Büyükköse [4] have obtained a lower bound for the smallest eigenvalue t_n of the $n \times n$ matrix $E_n^T E_n$, where the ij -entry of E_n is 1 if $j|i$ and 0 otherwise, i.e., $t_n \geq (n \sum_{k=1}^n \mu^2(k))^{-1}$. Indeed, this bound can be used instead of lower bounds including c_n for the smallest eigenvalues of GCD and related matrices defined on $S = \{1, 2, \dots, n\}$ in the literature, see [14,16,19,24,27]. After above studies on estimating the value of c_n , we naturally raise the following problem.

Problem 4.1. Can one improve the lower bounds mentioned above for c_n ?

In the review process, Professor Jorma K. Merikoski proposed a solution to Problem 4.1, which is presented in Appendix A.

Acknowledgements

The authors would like to thank Professor Ilker Inam, Professor Pentti Haukkanen and two anonymous reviewers for their valuable suggestions in improving the presentation

of our manuscript. We also would like to thank to Professor Jorma K. Merikoski for his solution to [Problem 4.1](#).

Appendix A. A solution to [Problem 4.1](#) (by Jorma K. Merikoski)

Recall that the matrix $Z_0^{-1} = (\zeta_{ij})$ satisfies

$$\zeta_{ii} = 1 + \sum_{k=i+1}^n f_{k-i}^2, \quad |\zeta_{ij}| = f_{|j-i|} + \sum_{k=j+1}^n f_{k-i} f_{k-j}, \quad i \neq j.$$

Also recall that the spectral radius of a nonnegative square matrix is less than or equal to the maximal row sum, and equality holds if and only if the matrix is irreducible and all row sums are equal (e.g., [\[9, Chapter 2, Theorem \(2.35\)\]](#)). Now

$$\begin{aligned} c_n^{-1} &= \rho(Z_0^{-1}) = \rho(|Z_0^{-1}|) < \max_i \sum_{j=1}^n |\zeta_{ij}| \\ &= \sum_{j=1}^n |\zeta_{1j}| \\ &= |\zeta_{11}| + \sum_{j=2}^n |\zeta_{1j}| \\ &= 1 + \sum_{k=2}^n f_{k-1}^2 + \sum_{j=2}^n \left(f_{j-1} + \sum_{k=j+1}^n f_{k-1} f_{k-j} \right) \\ &= 1 + \sum_{k=1}^{n-1} f_k + \sum_{j=1}^n \sum_{k=j+1}^n f_{k-1} f_{k-j} \\ &= 1 + S_1 + S_2. \end{aligned}$$

It is well-known [\[12, Eq. \(22\)\]](#) that

$$\sum_{k=1}^m f_k = f_{m+2} - 1; \tag{A.1}$$

so

$$S_1 = f_{n+1} - 1.$$

More work must be done with S_2 . First we have

$$\begin{aligned} S_2 &= (f_1^2 + f_2^2 + \cdots + f_{n-1}^2) + (f_2 f_1 + f_3 f_2 + \cdots + f_{n-1} f_{n-2}) + \cdots \\ &\quad + (f_j f_1 + f_{j+1} f_2 + \cdots + f_{n-1} f_{n-j}) + \cdots + (f_{n-2} f_1 + f_{n-1} f_2) + f_{n-1} f_1 \end{aligned}$$

$$\begin{aligned}
&= f_1(f_1 + \cdots + f_{n-1}) + f_2(f_2 + \cdots + f_{n-1}) + \cdots + f_j(f_j + \cdots + f_{n-1}) \\
&\quad + \cdots + f_{n-2}(f_{n-2} + f_{n-1}) + f_{n-1}^2 \\
&= \sum_{j=1}^{n-1} f_j \sum_{k=j}^{n-1} f_k.
\end{aligned}$$

By (A.1),

$$\sum_{k=j}^{n-1} f_k = \sum_{k=1}^{n-1} f_k - \sum_{k=1}^{j-1} f_k = (f_{n+1} - 1) - (f_{j+1} - 1) = f_{n+1} - f_{j+1},$$

which implies

$$S_2 = \sum_{j=1}^{n-1} f_j(f_{n+1} - f_{j+1}) = f_{n+1} \sum_{j=1}^{n-1} f_j - \sum_{j=1}^{n-1} f_j f_{j+1} = T_1 - T_2.$$

It remains to simplify T_1 and T_2 . By (A.1),

$$T_1 = f_{n+1}(f_{n+1} - 1).$$

It is well-known [29, Sequence A064831] that

$$\begin{aligned}
\sum_{j=1}^m f_j f_{j+1} &= f_{m+1}^2 - 1 \text{ if } n \text{ is even,} \\
\sum_{j=1}^m f_j f_{j+1} &= f_{m+1}^2 \text{ if } n \text{ is odd.}
\end{aligned}$$

Hence

$$\begin{aligned}
T_2 &= f_n^2 \text{ if } n \text{ is even,} \\
T_2 &= f_n^2 - 1 \text{ if } n \text{ is odd.}
\end{aligned}$$

In other words,

$$T_2 = f_n^2 - \eta_n,$$

where

$$\eta_n = \frac{1 - (-1)^n}{2}.$$

Now

$$\begin{aligned}
c_n^{-1} &< 1 + S_1 + T_1 - T_2 \\
&= 1 + (f_{n+1} - 1) + f_{n+1}(f_{n+1} - 1) - f_n^2 + \eta_n
\end{aligned}$$

$$\begin{aligned}
&= f_{n+1}^2 - f_n^2 + \eta_n \\
&= (f_{n+1} - f_n)(f_{n+1} + f_n) + \eta_n \\
&= f_{n-1}f_{n+2} + \eta_n.
\end{aligned}$$

It is well-known [8, Eq. (2)] that

$$f_{m-r}f_{m+s} - f_m f_{m+s-r} = (-1)^{m-r-1} f_r f_s.$$

Hence

$$f_{n-1}f_{n+2} = f_n f_{n+1} + (-1)^n,$$

and so

$$c_n^{-1} < f_n f_{n+1} + (-1)^n + \eta_n = f_n f_{n+1} + \theta_n,$$

where

$$\theta_n = \frac{1 + (-1)^n}{2}.$$

Therefore

$$c_n > (f_n f_{n+1} + \theta_n)^{-1} = \gamma_n,$$

which seems to be much better than Mattila's [25] bounds. For example, $c_6 = 0.0148$ (in three digits). The bound $\gamma_6 = 0.00952$ is rather good, while Mattila's better bound is 0.0000205.

References

- [1] E. Altınışık, On inverses of GCD matrices associated with multiplicative functions and a proof of the Hong–Loewy conjecture, *Linear Algebra Appl.* 430 (2009) 1313–1327.
- [2] E. Altınışık, On a conjecture on the smallest eigenvalues of some special positive definite matrices, in: 3rd International Conference on Applied Mathematics & Approximation Theory – AMAT 2015, 28–31 May 2015, Ankara, Turkey, 2015.
- [3] E. Altınışık, Ş. Büyükköse, A proof of a conjecture on monotonic behavior of the largest eigenvalue of a number-theoretic matrix, in: 12th International Conference of Numerical Analysis and Applied Mathematics, Rhodes, Greece, 2014.
- [4] E. Altınışık, Ş. Büyükköse, A proof of a conjecture on monotonic behavior of the smallest and the largest eigenvalue of a number-theoretic matrix, *Linear Algebra Appl.* 471 (2015) 141–149.
- [5] E. Altınışık, B.E. Sagan, N. Tuğlu, GCD matrices, posets, and nonintersecting paths, *Linear Multilinear Algebra* 53 (2) (2005) 75–84.
- [6] E. Altınışık, N. Tuğlu, P. Haukkanen, A note on bounds for norms of the reciprocal LCM matrix, *Math. Inequal. Appl.* 7 (4) (2004) 491–496.
- [7] F. Balatoni, On the eigenvalues of the matrix of the Smith determinant, *Mat. Lapok* 20 (1969) 397–403 (in Hungarian).
- [8] A.T. Benjamin, N.T. Cameron, J.J. Quinn, Fibonacci determinants – a combinatorial approach, *Fibonacci Quart.* 45 (2007) 39–55.

- [9] A. Berman, R.J. Plemmons, *Nonnegative Matrices in the Mathematical Sciences*, SIAM, 1994.
- [10] S. Beslin, S. Ligh, Greatest common divisor matrices, *Linear Algebra Appl.* 118 (1989) 69–76.
- [11] K. Bourque, S. Ligh, On GCD and LCM matrices, *Linear Algebra Appl.* 174 (1992) 65–74.
- [12] P. Chandra, E.W. Weisstein, Fibonacci number, from MathWorld – a Wolfram web resource, <http://mathworld.wolfram.com/FibonacciNumber.html>.
- [13] P. Haukkanen, J. Wang, J. Sillanpää, On Smith's determinant, *Linear Algebra Appl.* 258 (1997) 251–269.
- [14] S. Hong, Asymptotic behavior of largest eigenvalue of matrices associated with completely even functions (mod r), *Asian-Eur. J. Math.* 1 (2008) 225–235.
- [15] S. Hong, K.S. Enoch Lee, Asymptotic behavior of eigenvalues of reciprocal power LCM matrices, *Glasg. Math. J.* 50 (2008) 163–174.
- [16] S. Hong, R. Loewy, Asymptotic behavior of eigenvalues of greatest common divisor matrices, *Glasg. Math. J.* 46 (2004) 303–308.
- [17] S. Hong, R. Loewy, Asymptotic behavior of the smallest eigenvalue of matrices associated with completely even functions (mod r), *Int. J. Number Theory* 7 (2011) 1681–1704.
- [18] R. Horn, C.R. Johnson, *Matrix Analysis*, second ed., Cambridge University Press, Cambridge, London, 2013.
- [19] P. Ilmonen, P. Haukkanen, J.K. Merikoski, On eigenvalues of meet and join matrices associated with incidence functions, *Linear Algebra Appl.* 429 (2008) 859–874.
- [20] D. Kalman, A matrix proof of Newton's identities, *Math. Mag.* 73 (4) (2000) 859–874.
- [21] I. Korkee, P. Haukkanen, On meet and join matrices associated with incidence functions, *Linear Algebra Appl.* 372 (2003) 127–153.
- [22] T. Koshy, *Fibonacci and Lucas Numbers with Applications*, John Wiley, New York, 2001.
- [23] P. Lindqvist, K. Seip, Note on some greatest common divisor matrices, *Acta Arith.* 84 (2) (1998) 149–154.
- [24] M. Mattila, On the eigenvalues of combined meet and join matrices, *Linear Algebra Appl.* 466 (2015) 1–20.
- [25] M. Mattila, On the eigenvalues of combined meet and join matrices, *Linear Algebra Appl.* 466 (2015) 1–20.
- [26] M. Mattila, P. Haukkanen, On the eigenvalues of certain number-theoretic matrices, in: *International Conference in Number Theory and Applications*, 2012.
- [27] M. Mattila, P. Haukkanen, On the eigenvalues of certain number-theoretic matrices, *East-West J. Math.* 14 (2) (2012) 121–130.
- [28] M. Mattila, P. Haukkanen, On the positive definiteness and eigenvalues of meet and join matrices, *Discrete Math.* 326 (2014) 9–19.
- [29] N.J.A. Sloane, The on-line encyclopedia of integer sequences, <http://oeis.org/>.
- [30] H.J.S. Smith, On the value of a certain arithmetical determinant, *Proc. Lond. Math. Soc.* (1) 7 (1876) 208–212.
- [31] E. Süli, D. Mayers, *An Introduction to Numerical Analysis*, Cambridge University Press, Cambridge, London, 2003.
- [32] A. Wintner, Diophantine approximations and Hilbert's space, *Amer. J. Math.* 66 (1944) 564–578.